

APPLICATION OF NEUTRON ACTIVATION ANALYSIS FOR THE MEASUREMENT OF Br, Ca, Cl, K, Mg, Mn, and Na CONTENTS IN THE INTACT THYROID OF FEMALE

V. Zaichick¹, S. Zaichick^{1,2}

*¹ Medical Radiological Research Centre,
Koroleva str., 4, Obninsk, 249036, Russia,*

e-mail: vezai@obninsk.com

*² Feinberg School of Medicine, Northwestern University,
Chicago, IL 60611-4296, USA*

Introduction

A large proportion of the world and European populations has some evidence of thyroid dysfunction. For example, the prevalence of goiter in areas of severe iodine deficiency can be as high as 80%. In Germany, an area of relative iodine deficiency, thyroid nodules or goiter were found in 33% of working adults aged 18–65 years. Another problem is thyroid cancer. In the last decades, thyroid cancer incidence has continuously and sharply increased all over the world. Moreover, thyroid cancer mortality, in spite of earlier diagnosis and better treatment, has not decreased but is rather increasing. The reasons for this increase are not well understood, but some environmental carcinogens in the industrialized lifestyle may have specifically affected the thyroid. Among potential carcinogens, the increased iodine and some other chemical element intake is one of the most likely risk factors. Excessive accumulation or an imbalance of the chemical elements may disturb the cell functions and may result in cellular degeneration, death or, on the contrary, intensive uncontrolled proliferation, and malignancy. Questions on the role of chemical elements in etiology and pathogenesis of thyroid cancer are far from being answered. First of all, it is necessary to establish the normal level and age-related changes of chemical element contents in thyroidal parenchyma, and relationships of chemical elements in intact gland.

Aims of the study

This study aimed to perform a nondestructive method to evaluate the Br, Ca, Cl, K, Mg, Mn, and Na mass fraction in thyroidal parenchyma and present data on age-dependence and relationships of these element contents in intact thyroid of females. To that end, we determined Br, Ca, Cl, K, Mg, Mn, and Na mass fraction in intact thyroid glands using an instrumental neutron activation analysis with high resolution spectrometry of short-lived radionuclides..

All studies were approved by the Ethical Committees of the Medical Radiological Research Centre, Obninsk.

Materials

Samples of the human thyroid were obtained from randomly selected autopsy specimens of 33 females (European-Caucasian) aged 3.5 to 87 years. All the deceased were citizens of Obninsk and had undergone routine autopsy at the Forensic Medicine Department of City Hospital, Obninsk. Age ranges for subjects were divided into two age groups, with group 1, 3.5-40 years (30.9 ± 3.1 years, $M \pm SEM$, $n=11$) and group 2, 41–87 years (66.3 ± 2.7 years, $M \pm SEM$, $n=22$). These groups were selected to reflect the condition of thyroid tissue in the children, teenagers, young adults and first period of adult life (group 1) and in the second period of adult life as well as in old age (group 2). The available clinical data were reviewed for each subject. None of the subjects had a history of an intersex condition, endocrine disorder, or other chronic disease that could affect the normal development of the thyroid. None of the subjects were receiving medications or used any supplements known to affect thyroid chemical element contents. The typical causes of sudden death of most of these subjects included trauma or suicide and also acute illness (cardiac insufficiency, stroke, embolism of pulmonary artery, alcohol poisoning).

All right lobes of thyroid glands were divided into two portions using a titanium scalpel. One tissue portion was reviewed by an anatomical pathologist while the other was used for the chemical element content determination. A histological examination was used to control the age norm conformity as well as the unavailability of microadenomatosis and latent cancer.

Methods

All samples were freeze-dried and homogenized. The pounded sample weighing about 100 mg was used for measurement by INAA-SLR. The samples for INAA-SLR were sealed separately in thin polyethylene films washed beforehand with acetone and rectified alcohol. The sealed samples were placed in labeled polyethylene ampoules.

The content of Br, Ca, Cl, I, K, Mg, Mn, and Na were determined by INAA-SLR using a horizontal channel equipped with the pneumatic rabbit system of the WWR-c research nuclear reactor. The neutron flux in the channel was $1.7 \times 10^{13} \text{ n cm}^{-2} \text{ s}^{-1}$. Ampoules with thyroid tissue samples, biological synthetic standards, intralaboratory-made standards, and certified reference material (CRM) IAEA H-4 (animal muscle) were put into polyethylene rabbits and then irradiated separately for 180 s. Copper foils were used to assess neutron flux.

The measurement of each sample was made twice, 1 and 120 min after irradiation. The duration of the first and second measurements was 10 and 20 min, respectively. A coaxial 98-cm³ Ge (Li) detector and a spectrometric unit (NUC 8100), including a PC-coupled multichannel analyzer, were used for measurements. The spectrometric unit provided 2.9-keV resolution at the ⁶⁰Co 1,332-keV line. Details of used nuclear reactions, radionuclides, and gamma-energies were presented in our earlier publications concerning the INAA chemical element contents in human scalp hair.

Results and Discussion

Good agreement of the Br, Ca, Cl, I, K, Mg, Mn, and Na contents analyzed by INAA-SLR with the certified data of CRM IAEA H-4 (Table 1) indicates an acceptable accuracy of the results obtained in the study of chemical elements of the thyroid presented in Tables 2–5.

Table 1. INAA-SLR data of chemical element contents in the IAEA H-4 (animal muscle) reference material compared to certified values (mg/kg, dry mass basis)

Element	Certified values			Type	This work results
	Mean	95% confidence interval			
Br	4.1	3.5 – 4.7		N	5.0±0.9
Ca	188	163 – 213		N	238±59
Cl	1890	1810 – 1970		N	1950±230
I	0.08	-		N	<1.0
K	15800	15300 – 16400		C	16200±3800
Mg	1050	990 – 1110		C	1100±190
Mn	0.52	0.48 – 0.55		C	0.55±0.11
Na	2060	1930 – 2180		C	2190±140

Mean - arithmetical mean, SD - standard deviation, C - certified values, N - non-certified values.

Table 2 represents certain statistical parameters (arithmetic mean, standard deviation, standard error of mean, minimal and maximal values, median, percentiles with 0.025 and 0.975 levels) of the Br, Ca, Cl, I, K, Mg, Mn, and Na mass fractions in intact (normal) thyroid of females.

Table 2.. Some statistical parameters of Br, Ca, Cl, I, K, Mg, Mn, and Na mass fraction (mg/kg, dry mass basis) in intact thyroid of female

Gender	Element	Mean	SD	SEM	Min	Max	Median	P 0.025	P 0.975
Females n=33	Br	22.4	16.1	3.2	5.00	66.9	16.3	5.00	59.2
	Ca	1663	570	198	461	3640	1170	670	3600
	Cl	3317	1480	290	1200	6000	3375	1388	5906
	I	1956	1199	219	114	5061	1562	309	4662
	K	5395	3245	723	1740	13700	4835	2120	13230
	Mg	212	97	24	66.0	364	215	67.5	356
	Mn	1.50	0.84	0.22	0.550	4.18	1.37	0.603	3.41
	Na	6421	1721	320	3800	10450	6700	4122	9924

Mean – arithmetic mean, SD – standard deviation, SEM – standard error of mean, Min – minimum value, Max – maximum value, P 0.025 – percentile with 0.025 level, P 0.975 – percentile with 0.975 level.

The obtained means for Br, Ca, Cl, I, K, Mg, Mn, and Na mass fraction, as shown in Table 3, agree well with the medians of mean values cited by other researches for the human thyroid, including samples received from persons who died from different non-thyroid diseases [25-36]. A number of values for mass fractions were not expressed on a dry mass basis by the authors of the cited references. However, we calculated these values using published data for water (75%) and ash (4.16% on dry mass basis) contents in thyroid of adults.

Table 3. Median, minimum and maximum value of means Br, Ca, Cl, I, K, Mg, Mn, and Na contents in normal thyroid according to data from the literature in comparison with our results (mg/kg, dry mass basis)

Element	Published data [Reference]			This work
	Median of means (n)*	Minimum of means M or M±SD, (n)**	Maximum of means M or M±SD, (n)**	
Br	18.1 (11)	5.12 (44) [25]	284±44 (14) [26]	22±16
Ca	1600 (17)	840±240 (10) [27]	3800±320 (29) [27]	1663±570
Cl	6800 (5)	804±80 (4) [28]	8000 (-) [29]	3317±1480
I	1888 (95)	159±8 (23) [30]	5772±2708 (50) [31]	1956±1199
K	4400 (17)	46.4±4.8 (4) [28]	6090 (17) [32]	5395±3245
Mg	390 (16)	3.5 (-) [33]	840±400 (14) [34]	212±97
Mn	1.82 (36)	0.44±11 (12) [35]	69.2±7.2 (4) [28]	1.50±0.84
Na	8000 (9)	438 (-) [36]	10000±5000 (11) [34]	6421±1721

M – arithmetic mean, SD – standard deviation, (n)* – number of all references, (n)** – number of samples.

To estimate the effect of age on the chemical element contents we examined two age groups, described above (Table 4). Moreover, a correlation between age and elemental mass fraction was calculated (Table 5).

Table 4. Differences between mean values ($M \pm SEM$) of Br, Ca, Cl, I, K, Mg, Mn, and Na mass fraction (mg/kg, dry mass basis) in normal female thyroid of two age groups (AG)

Element	Female thyroid tissue				Ratio
	AG1 3.5-40 years n=11	AG2 41-87 years n=22	t-test $p \leq$	U-test p	AG2 to AG1
Br	13.3±2.5	26.8±4.3	0.0118	≤0.01	2.02
Ca	1052±65	2029±276	0.0034	≤0.01	1.93
Cl	4109±544	2965±318	0.0947	≤0.05	0.72
I	1876±346	2002±288	0.782	>0.05	1.07
K	5379±1101	5408±1013	0.984	>0.05	1.01
Mg	212±39	212±31	0.994	>0.05	1.00
Mn	1.43±0.13	1.57±0.46	0.772	>0.05	1.10
Na	5969±458	6025±414	0.300	>0.05	1.01

M – arithmetic mean, SEM – standard error of mean, *t*-test - Student's *t*-test, U-test - Wilcoxon-Mann-Whitney *U*-test, Statistically significant values are in **bold**.


Table 5. Correlations between age and chemical element mass fractions in the intact thyroid of female (*r* – coefficient of correlation)

Element	Br	Ca	Cl	I	K	Mg	Mn	Na
Age	0.181	0.369 ^a	-0.425 ^a	0.084	-0.051	-0.197	-0.036	0.177

Statistically significant values: ^a $p \leq 0.05$.

A strongly pronounced tendency of age-related increase in Br and Ca mass fraction was observed in thyroid when two age groups were compared (Table 4). In second group of females with mean age 66.3 years the mean Br and Ca mass fraction in thyroids was almost 2 times higher than in thyroids of the first age group (mean age 30.9 years). However, the correlation between age and elemental mass fractions found a significant direct dependence for Ca and inverse for Cl (Table 5). Why was the direct correlation between age and Br mass fraction not found?

Fig. 1. Data sets of individual Br, Ca, and Cl mass fraction values in intact thyroid of females and their trend lines



A significant direct correlation between the Br-K, Br-Mn, Cl-Mg, Cl-Na, I-Na, K-Mg, and Mn-Na mass fractions as well as an inverse correlation between Ca-Cl and Na-Mn mass fractions was seen in female thyroid. No correlation was demonstrated between any other chemical elements (Table 5). If some correlations between the elements were predictable (e.g., Na-Cl), the interpretation of other observed relationships would require further study.

Table 6. Intercorrelations of the chemical element mass fractions in the intact thyroid of female (*r* – coefficient of correlation)

Element	Br	Ca	Cl	I	K	Mg	Mn	Na
Br	1.00	0.246	0.073	-0.053	0.398 ^a	-0.179	0.727 ^c	0.179
Ca	0.246	1.00	-0.658 ^c	0.132	0.125	0.021	0.341	-0.184
Cl	0.073	-0.658 ^c	1.00	0.159	-0.367	0.563 ^b	-0.207	0.389 ^a
I	-0.053	0.132	0.159	1.00	-0.319	0.191	-0.090	0.446 ^a
K	0.398 ^a	0.125	-0.367	-0.319	1.00	0.774 ^c	-0.194	-0.152
Mg	-0.179	0.021	0.563 ^b	0.191	0.774 ^c	1.00	-0.266	0.313
Mn	0.727 ^c	0.341	-0.207	-0.090	-0.194	-0.266	1.00	-0.410 ^a
Na	0.179	-0.184	0.389 ^a	0.446 ^a	-0.152	0.313	0.410 ^a	1.00

Statistically significant values: ^a $p \leq 0.05$, ^b $p \leq 0.01$, ^c $p \leq 0.001$.

The Br is one of the most abundant and ubiquitous of the recognized trace elements in the biosphere. Inorganic bromide is the ionic form of bromine which exerts therapeutic as well as toxic effects. An enhanced intake of bromide could interfere with the metabolism of iodine at the whole-body level. In the thyroid gland the biological behavior of bromide is more similar to the biological behavior of iodide. Therefore, a goitrogenic effect of excessive bromide level in the thyroid of old females may be assumed.

In addition to the elevated Br level, an age-related increase and excess in Ca mass fractions in thyroid tissue may contribute to harmful effects on the gland. There are good reasons for such speculations since many reviews and numerous papers raise the concern about role of Ca in the prostate, breast, lung and other organ malignant transformation. Calcium ions Ca^{2+} are central to both cell proliferation and cell death. Changes in cytosolic Ca^{2+} trigger events critical for tumorigenesis, such as cellular motility, proliferation and apoptosis. An increased growth rate of cells is correlated with an increase in the intracellular calcium pool content. Moreover, increases in cytosolic free Ca^{2+} represent a ubiquitous signalling mechanism that controls a variety of cellular processes, including not only proliferation, but also cell metabolism and gene transcription. Indeed, an increased level of Ca content in the thyroid tissue of old females reflects an increase in the intracellular calcium pool. Thus, an increase of Ca content in tissue and organs with age is a key feature in etiology of many benign and malignant tumors, including thyroid goiter and cancer.

Conclusions

The instrumental neutron activation analysis with high resolution spectrometry of short-lived radionuclides is a useful analytical tool for the non-destructive determination of chemical element content in the thyroid tissue samples. This method allows determine means for Br, Ca, Cl, I, K, Mg, Mn, and Na (8 chemical elements).

Our data reveal that there is a strongly pronounced tendency of increase in Br and Ca mass fraction as well as of decrease in Cl mass fraction in the normal thyroid of female during a lifespan. Therefore, a goitrogenic and tumorogenic effect of excessive Br and Ca level in the thyroid of old females may be assumed.

All the deceased were citizens of Obninsk. Obninsk is the small nonindustrial city not far from Moscow in unpolluted area. None of those who died a sudden death had suffered from any systematic or chronic disorders before. The normal state of thyroid was confirmed by morphological study. Thus, our data for Br, Ca, Cl, I, K, Mg, Mn, and Na mass fractions in intact thyroid may serve as indicative normal values for females of urban population of the Russian Central European region.