

# RELATIONSHIP BETWEEN Ca, Cl, K, Mg, Mn, Na, P, AND Sr CONTENTS IN THE INTACT ROOTS OF MALE TEETH INVESTIGATED BY NEUTRON ACTIVATION ANALYSIS

V. Zaichick<sup>1</sup>, S. Zaichick<sup>1,2</sup>

<sup>1</sup>*Medical Radiological Research Centre, Korolyev St., 4, Obninsk, 249036, Russia,  
e-mail: [vezai@obninsk.com](mailto:vezai@obninsk.com)*

<sup>2</sup>*Feinberg School of Medicine, Northwestern University, Chicago, IL 60611-4296, USA*

## Introduction

To use chemical element composition as estimation of bone and teeth health in stomatological, geographical, environmental and occupational medicine, paleoanthropology, and other directions, it is necessary to know normal levels and age- and gender-related changes of chemical element contents and their ratios.<sup>[1-22]</sup>

In our previous studies it was shown that samples are contaminated by trace elements from stainless steel tools during the sample preparation.<sup>[23]</sup> There is evidence that some chemical elements are lost if tooth samples are treated with solvents in order to remove the organic matrix, and are then ashed and acid digested.<sup>[24, 25]</sup>

This work had three aims. The first was to determine the Ca, Cl, K, Mg, Mn, Na, P, and Sr mass fractions in the intact roots of male teeth by instrumental neutron activation analysis with high resolution spectrometry of short-lived radionuclides (INAA-SLR) and to calculate some statistical parameters of Cl/Ca, K/Ca, Mg/Ca, Mn/Ca, Na/Ca, P/Ca, Sr/Ca, Ca/P, Cl/P, K/P, Mg/P, Mn/P, Na/P, Sr/P, Ca/Mg, Cl/Mg, Mn/Mg, Na/Mg, P/Mg, Sr/Mg, Ca/Cl, K/Cl, Mg/Cl, Mn/Cl, Na/Cl, P/Cl, Sr/Cl, Ca/K, Cl/K, Mg/K, Mn/K, Na/K, P/K, Sr/K, Ca/Na, Cl/Na, K/Na, Mg/Na, Mn/Na, P/Na, and Sr/Na mass fraction ratios. The second aim was to evaluate the effect of age on mean values of ratios of chemical element mass fractions. The third aim was to estimate the inter correlations between Ca, Cl, K, Mg, Mn, Na, P, and Sr mass fractions in the intact roots of male teeth.

All studies were approved by the Ethical Committee of the Medical Radiological Research Center, Obninsk.

## Experimental

Tooth root samples were obtained at postmortems from intact cadavers (46 male, 16–58 years old) within 24 h of death. Samples were freeze dried until constant mass was obtained. All subjects died mainly due to trauma. A titanium tool was used to cut and to scrub samples as well as to cut teeth roots. The samples weighing about 100 mg for INAA-SLR were sealed separately in thin polyethylene film washed beforehand with acetone and rectified alcohol. The sealed samples were placed in labeled polyethylene ampoules.

To determine contents of the elements by comparison with a known standard, biological synthetic standards (BSS) prepared from phenol–formaldehyde resins and chemically pure compounds were used.<sup>[26]</sup> Ten certified reference materials (CRM) IAEA H-5 (Animal Bone)

and standard reference materials (SRM) NIST 1486 (Bone Meal) sub-samples weighing about 50–100mg were analyzed in the same conditions as teeth samples to estimate the precision and accuracy of results.

The mass fractions of Ca, Cl, K, Mg, Mn, Na, P, and Sr were determined by INAA-SLR using a horizontal channel equipped with the pneumatic rabbit system of the WWR-c research nuclear reactor. The information of used nuclear reactions, radionuclides, gamma-energies, neutron flux, spectrometer and other details of the analysis including the quality control of results were reported by us before.<sup>[6, 26]</sup>

A dedicated computer program of NAA mode optimization was used.<sup>[27]</sup> Using the Microsoft Office Excel software, the summary of statistics were calculated for different ratios of chemical element mass fractions. The reliability of difference in the results between two age groups of females was evaluated by Student's *t*-test. The Pearson's correlation analysis was used to identify relationships between elements.

## Results and discussion

Table 1 represents certain statistical parameters (arithmetic mean, standard deviation, standard error of mean, minimal and maximal values, median, percentiles with 0.025 and 0.975 levels) of 42 different ratios of Ca, Cl, K, Mg, Mn, Na, P, and Sr mass fractions in the intact roots of male teeth.

The obtained values for Ca/P ratio, as shown in Table 1, agree well with median of means cited by other researchers for the teeth enamel.<sup>[28, 29]</sup> No published data referring to ratios of Ca/P or other chemical element mass fractions in the intact roots of male teeth was found.

To estimate the effect of age on the chemical element ratios in the intact roots of male teeth we examined two age groups: one comprised a younger group with ages from 15 to 35 years and the other comprised older people with ages ranging from 36 to 55 years (Table 2). In the intact roots of male teeth it was not found the statistically significant age-related changes of the mass fraction ratios of chemical elements investigated in the study.

No published data referring to age-related differences of ratios of chemical element mass fractions in the intact roots of male teeth were found.

Table 3 depicts the inter-correlation calculations including all chemical elements identified by us. For example, the positive inter-correlations of Ca mass fractions with P ( $p < 0.001$ ), Cl ( $p < 0.001$ ), and Mn ( $p < 0.01$ ) mass fractions were found in male teeth roots. If some correlations between the elements were predictable (e.g., Ca–P), the interpretation of other observed relationships requires further study for a more complete understanding.

## Conclusions

INAA-SLR is a satisfactory analytical tool for the non-destructive precise determination of mass fraction of 8 chemical elements (Ca, Cl, K, Mg, Mn, Na, P, and Sr) in the root samples of permanent teeth. It was not found the statistically significant changes of chemical element mass fraction ratios in the intact roots of male teeth with age. All the deceased were citizens of Obninsk, a small city of non-industrial region near Moscow. None of those who died a sudden death had suffered from any systematic or chronic disorders before. Thus, our data for element mass fraction ratios in the intact roots of male teeth may serve as indicative normal values for residents of the Central European region of Russia.

**Table 1.** Some statistical parameters of 42 different ratios of Ca, Cl, K, Mg, Mn, Na, P, and Sr mass fractions in the intact roots of male teeth

Ratio	M	SD	SEM	Min	Max	Med	P0.025	P0.975
(Cl/Ca)×10 <sup>3</sup>	3.83	2.34	0.36	0.880	14.2	3.43	1.27	7.63
(K/Ca) ×10 <sup>3</sup>	3.46	1.13	0.19	0.587	6.69	3.10	1.59	6.11
(Mg/Ca) ×10 <sup>2</sup>	2.97	0.83	0.13	0.692	4.55	3.09	1.42	4.29
(Mn/Ca) ×10 <sup>6</sup>	5.20	3.25	0.51	1.25	15.8	3.99	1.49	11.3
(Na/Ca) ×10 <sup>2</sup>	2.59	1.13	0.18	1.69	7.98	2.28	1.74	5.90
P/Ca	0.543	0.056	0.009	0.427	0.714	0.541	0.454	0.635
(Sr/Ca) ×10 <sup>3</sup>	1.79	1.00	0.15	0.231	5.36	1.61	0.511	3.59
Ca/P	1.86	0.19	0.03	1.40	2.34	1.85	1.58	2.20
(Cl/P) ×10 <sup>2</sup>	0.713	0.430	0.066	0.164	2.52	0.658	0.233	1.35
(K/P) ×10 <sup>2</sup>	0.626	0.202	0.034	0.129	1.24	0.575	0.349	1.05
(Mg/P) ×10 <sup>2</sup>	5.54	1.71	0.27	1.22	8.92	5.46	2.62	8.57
(Mn/P) ×10 <sup>6</sup>	9.56	6.04	0.96	2.53	29.5	6.90	2.78	21.9
(Na/P) ×10 <sup>2</sup>	4.79	1.98	0.31	2.81	13.5	4.30	3.07	10.6
(Sr/P) ×10 <sup>3</sup>	3.30	1.81	0.27	0.467	8.97	3.03	0.845	7.26
Ca/Mg	38.1	19.9	3.10	22.0	145	32.4	23.3	70.6
Cl/Mg	0.158	0.180	0.028	0.0254	1.11	0.114	0.0384	0.482
K/Mg	0.136	0.072	0.012	0.0179	0.419	0.126	0.0455	0.273
(Mn/Mg) ×10 <sup>4</sup>	2.04	1.51	0.24	0.320	6.90	1.40	0.428	5.60
Na/Mg	0.952	0.503	0.079	0.457	3.11	0.789	0.504	2.58
P/Mg	20.8	11.5	1.8	11.2	82.0	18.3	11.7	38.2
Sr/Mg	0.0680	0.0487	0.0070	0.0059	0.275	0.0592	0.0152	0.159
Ca/Cl	355	210	32	70.7	1136	292	131	791
K/Cl	1.30	0.93	0.16	0.213	4.41	1.03	0.354	3.68
Mg/Cl	11.0	7.8	1.2	0.904	39.4	8.75	2.07	26.0
(Mn/Cl) ×10 <sup>4</sup>	19.6	27.9	4.4	2.32	180	13.9	3.75	53.3
Na/Cl	9.41	7.82	1.21	1.69	47.1	7.07	2.34	24.5
P/Cl	194	119	18	39.5	610	152	74.2	430
Sr/Cl	0.645	0.658	0.099	0.0698	3.98	0.447	0.0757	1.65
Ca/K	344	250	42	149	1704	322	164	737
Cl/K	1.20	0.89	0.15	0.227	4.69	0.973	0.274	2.95
Mg/K	9.94	8.77	1.51	2.39	55.8	7.95	3.82	25.8
(Mn/K) ×10 <sup>4</sup>	16.2	11.3	1.9	7.02	49.4	10.9	7.40	42.1
Na/K	8.69	7.39	1.25	4.24	48.6	6.67	4.83	20.0
P/K	185	111	19	80.7	774	174	95.4	335
Sr/K	0.529	0.271	0.042	0.172	1.31	0.468	0.187	1.25
Ca/Na	42.4	9.8	1.5	12.5	59.1	43.9	17.0	57.6
Cl/Na	0.165	0.114	0.018	0.0212	0.593	0.141	0.0409	0.428
K/Na	0.141	0.043	0.007	0.0206	0.236	0.150	0.0602	0.208
Mg/Na	1.23	0.42	0.07	0.322	2.19	1.27	0.388	1.98
(Mn/Na) ×10 <sup>4</sup>	2.24	1.66	0.26	0.415	7.33	1.68	0.520	6.37
P/Na	23.0	5.9	0.9	7.38	35.6	23.3	9.42	32.6
Sr/Na	0.746	0.405	0.061	0.0766	1.67	0.641	0.231	1.61

*M* arithmetic mean, *SD* standard deviation, *SEM* standard error of mean, *Min* minimum value, *Max* maximum value, *Med* median, *P0.025* percentile with 0.025 level, *P0.975* percentile with 0.975 level

**Table 2.** Effect of age on mean values (M±SEM) of ratios of chemical element in the intact roots of male teeth (Student's *t*-test)

Ratio	Age			Ratio M <sub>2</sub> /M <sub>1</sub>
	15-35 year (M <sub>1</sub> )	36-55 year (M <sub>2</sub> )	<i>p</i> ≤	
(Cl/Ca)×10 <sup>3</sup>	3.99±0.41	3.70±0.56	0.67	0.93
(K/Ca) ×10 <sup>3</sup>	3.41±0.34	3.50±0.22	0.83	1.03
(Mg/Ca) ×10 <sup>2</sup>	2.88±0.19	3.03±0.18	0.55	1.05
(Mn/Ca) ×10 <sup>6</sup>	4.74±0.62	5.57±0.79	0.41	1.18
(Na/Ca) ×10 <sup>2</sup>	2.79±0.38	2.44±0.12	0.37	0.87
P/Ca	0.534±0.012	0.550±0.012	0.37	1.03
(Sr/Ca) ×10 <sup>3</sup>	1.78±0.18	1.80±0.24	0.94	1.01
Ca/P	1.89±0.05	1.84±0.04	0.41	0.97
(Cl/P) ×10 <sup>2</sup>	0.745±0.072	0.690±0.104	0.67	0.93
(K/P) ×10 <sup>2</sup>	0.625±0.056	0.627±0.043	0.98	1.00
(Mg/P) ×10 <sup>2</sup>	5.46±0.39	5.59±0.37	0.81	1.02
(Mn/P) ×10 <sup>6</sup>	6.16±1.20	10.2±1.5	0.46	1.66
(Na/P) ×10 <sup>2</sup>	5.23±0.65	4.46±0.22	0.27	0.85
(Sr/P) ×10 <sup>3</sup>	3.30±0.34	3.30±0.43	0.99	1.00
Ca/Mg	37.5±2.8	38.5±5.0	0.86	1.03
Cl/Mg	0.157±0.027	0.159±0.045	0.97	1.01
K/Mg	0.131±0.015	0.139±0.019	0.75	1.06
(Mn/Mg) ×10 <sup>4</sup>	2.00±0.40	2.07±0.31	0.88	1.04
Na/Mg	1.02±0.14	0.905±0.090	0.50	0.89
P/Mg	20.1±1.7	21.3±2.9	0.72	1.06
Sr/Mg	0.065±0.008	0.071±0.012	0.70	1.09
Ca/Cl	320±44	382±46	0.34	1.19
K/Cl	1.12±0.23	1.45±0.21	0.30	1.29
Mg/Cl	9.79±1.66	11.9±1.7	0.38	1.22
(Mn/Cl) ×10 <sup>4</sup>	13.4±1.8	24.7±7.8	0.17	1.84
Na/Cl	9.75±2.5	9.15±1.05	0.83	0.94
P/Cl	169±22	214±27	0.21	1.27
Sr/Cl	0.551±0.096	0.731±0.169	0.36	1.33
Ca/K	394±91	301±14	0.32	0.76
Cl/K	1.34±0.17	1.08±0.24	0.39	0.81
Mg/K	11.6±3.3	8.67±0.71	0.41	0.75
(Mn/K) ×10 <sup>4</sup>	16.2±3.3	16.2±2.2	0.99	1.00
Na/K	10.5±2.7	7.15±0.35	0.23	0.68
P/K	202±40	170±9	0.44	0.84
Sr/K	0.541±0.058	0.515±0.064	0.76	0.95
Ca/Na	41.8±2.9	42.8±1.6	0.75	1.02
Cl/Na	0.173±0.023	0.160±0.026	0.71	0.92
K/Na	0.136±0.014	0.145±0.007	0.56	1.07
Mg/Na	1.15±0.09	1.29±0.09	0.28	1.12
(Mn/Na) ×10 <sup>4</sup>	2.06±0.38	2.38±0.37	0.54	1.16
P/Na	22.3±1.7	23.5±1.0	0.55	1.05
Sr/Na	0.736±0.085	0.755±0.089	0.88	1.03

*M* arithmetic mean, *SEM* standard error of mean, *bold* statistically significant

**Table 3.** Intercorrelations of the chemical element mass fractions in the intact roots of male teeth ( $r$  – coefficient of correlation)

Element	Ca	Cl	K	Mg	Mn	Na	P	Sr
Ca	<b>1.00</b>	0.594 <sup>c</sup>	0.126	0.205	0.393 <sup>b</sup>	-0.049	0.903 <sup>c</sup>	0.137
Cl	0.594 <sup>c</sup>	<b>1.00</b>	0.279	0.022	0.260	0.091	0.516 <sup>c</sup>	0.058
K	0.126	0.279	<b>1.00</b>	-0.175	0.240	-0.088	0.205	0.209
Mg	0.205	0.022	-0.175	<b>1.00</b>	0.002	-0.018	0.138	-0.142
Mn	0.393 <sup>b</sup>	0.260	0.240	0.002	<b>1.00</b>	-0.112	0.368 <sup>a</sup>	0.223
Na	-0.049	0.091	-0.088	-0.018	-0.112	<b>1.00</b>	-0.018	-0.032
P	0.903 <sup>c</sup>	0.516 <sup>c</sup>	0.205	0.138	0.368 <sup>a</sup>	-0.018	<b>1.00</b>	0.111
Sr	0.137	0.058	0.209	-0.142	0.223	-0.032	0.111	<b>1.00</b>

Statistically significant values: <sup>a</sup> $p \leq 0.05$ , <sup>b</sup> $p \leq 0.01$ , <sup>c</sup> $p \leq 0.001$ .

### References

1. Zaichick V. (1993) The *in vivo* neutron activation analysis of calcium in the skeleton of normal subjects, with hypokinesia and bone diseases. *J. Radioanal. Nucl. Chem.* **169**(2): 307–316.
2. Zaichick V. (1994) Instrumental activation and X-ray fluorescent analysis of human bones in health and disease. *J. Radioanal. Nucl. Chem.* **179**(2):295–303.
3. Zaichick V., Ovchjarenko N.N. (1996) *In vivo* X-ray fluorescent analysis of Ca, Zn, Sr, and Pb in frontal tooth enamel. *J. Trace and Microprobe Techniques* **14**(1):143–152.
4. Zaichick V., Morukov B. (1998) *In vivo* bone mineral studies on volunteers during a 370-day antiorthostatic hypokinesia test. *J. Appl. Radiat. Isot.* **49**(5/6): 691–694.
5. Zaichick V., Ovchjarenko N., Zaichick S. (1999) *In vivo* energy dispersive x-ray fluorescence for measuring the content of essential and toxic trace elements in teeth. *J. Appl. Radiat. Isot.* **50**(2):283–293.
6. Zaichick V., Dyatlov A., Zaichick S. (2000) INAA application in the age dynamics assessment of major, minor, and trace elements in the human rib. *J. Radioanal. Nucl. Chem.* **244**:189–193.
7. Zaichick V., Tzaphlidou M. (2002) Determination of calcium, phosphorus, and the calcium/phosphorus ratio in cortical bone from the human femoral neck by neutron activation analysis. *J. Appl. Radiat. Isot.* **56**:781–786.
8. Tzaphlidou M., Zaichick V. (2002) Neutron activation analysis of calcium/phosphorus ratio in rib bone of healthy humans. *J. Appl. Radiat. Isot.* **57**: 779–783.
9. Tzaphlidou M., Zaichick V. (2003) Calcium, phosphorus, calcium-phosphorus ratio in rib bone of healthy humans. *Biol. Trace Elem. Res.* **93**:63–74.
10. Zaichick V., Tzaphlidou M. (2003) Calcium and phosphorus concentrations and calcium/phosphorus ratio in trabecular bone from femoral neck of healthy humans as determined by neutron activation analysis. *J. Appl. Radiat. Isot.* **58**:623–627.
11. Zaichick V. (2004) INAA application in the age dynamics assessment of Ca, Cl, K, Mg, Mn, Na, P, and Sr contents in the cortical bone of human femoral neck. *J. Radioanal. Nucl. Chem.* **259**:351–354.
12. Tzaphlidou M., Zaichick V. (2004) Sex and age related Ca/P ratio in cortical bone of iliac crest of healthy humans. *J. Radioanal. Nucl. Chem.* **259**(2): 347–349.
13. Zaichick V. (2006) NAA of Ca, Cl, K, Mg, Mn, Na, P, and Sr contents in the human cortical and trabecular bone. *J. Radioanal. Nucl. Chem.* **269**:653–659.

14. Zaichick V. (2007) INAA application in the assessment of selected elements in cancellous bone of human iliac crest. *J. Radioanal. Nucl. Chem.* **271**:573–576.
15. Zaichick V., Zaichick S. (2009) Instrumental neutron activation analysis of trace element contents in the rib bone of healthy men. *J. Radioanal. Nucl. Chem.* **281**:47–52.
16. Zaichick V. (2009) Neutron activation analysis of Ca, Cl, K, Mg, Mn, Na, P, and Sr contents in the crowns of human permanent teeth. *J. Radioanal. Nucl. Chem.* **281**(1):41–45.
17. Zaichick S., Zaichick V. (2010) The effect of age and gender on 38 chemical element contents in human iliac crest investigated by instrumental neutron activation analysis. *J. Trace Elements. Med. Biol.* **24**: 1–6.
18. Zaichick S., Zaichick V. (2010) The effect of age and gender on 38 chemical element contents in human femoral neck investigated by instrumental neutron activation analysis”. *Biol. Trace Elem. Res.* **137**: 1–12.
19. Zaichick S., Zaichick V. (2012) Neutron activation analysis of Ca, Cl, Mg, Na, and P content in human bone affected by osteomyelitis or osteogenic sarcoma. *J. Radioanal. Nucl. Chem.* **293**: 241–246.
20. Zaichick V. (2013) Chemical elements of human bone tissue investigated by nuclear analytical and related methods. *Biol. Trace Elem. Res.* **153**: 84–99.
21. Zaichick V. (2013) Data for the Reference Man: skeleton content of chemical elements. *Radiat. Environ. Biophys.* **52**(1): 65–85.
22. Zaichick V., Zaichick S. (2014) The Ca, Cl, Mg, Na, and P mass fractions in human bone affected by Ewing’s sarcoma. *Biol. Trace Elem. Res.* **159**(1-3):32–38.
23. Zaichick V., Zaichick S. (1996) Instrumental effect on the contamination of biomedical samples in the course of sampling. *J. Anal. Chem.* **51**(12):1200–1205.
24. Zaichick V. (1997) Sampling, sample storage and preparation of biomaterials for INAA in clinical medicine, occupational and environmental health. In: *Harmonization of Health-Related Environmental Measurements Using Nuclear and Isotopic Techniques*. IAEA, Vienna, pp 123–133.
25. Zaichick V. (2004) Losses of chemical elements in biological samples under the dry ashing process. *Trace Elements in Medicine* **5**(3): 17–22.
26. Zaichick V. (1995) Application of synthetic reference materials in the Medical Radiological Research Centre. *Fresenius J. Anal. Chem.* **352**:219–223.
27. Korelo A.M., Zaichick V. (1993) Software to optimize the multielement INAA of medical and environmental samples. In: *Activation Analysis in Environment Protection*. Joint Institute on Nuclear Research, Dubna, Russia, pp 326–332.
28. Antunes A., Salvador V.L.R., Scapin M.A., de Rossi W., Zezell D.M. (2005) Nanosecond Nd: YAG laser on dental enamel: compositional analysis by X-ray fluorescence. *Laser Phys. Lett.* **2**(6): 318–323.
29. Liu H., Chao J., Chuang C., Chiu H., Yang C., Sun Y. (2013) Study of P, Ca, Sr, Ba and Pb levels in enamel and dentine of human third molars for environmental and archaeological research. *Advances in Anthropology* **3**: 71–77.